

Powerful MRI could yield new way to monitor MS progression


University of Alberta research study funded in part by MS Society of Canada

Researchers at the University of Alberta have discovered a new way to track the progression of MS by using a powerful, triple strength MRI to track increasing levels of iron found in brain tissue.

The researchers discovered that iron levels in MS patients are increasing in grey matter areas of the brain that are responsible for relaying messages. High iron levels in a specific “relay area” were noted in patients who had physical disabilities associated with MS. Iron is very important for normal function of the brain and the amount of iron is a tightly controlled system by the brain tissue. The discovery suggests there is a problem with the control system. Too much iron can be toxic to brain cells and high levels of iron in the brain have been associated with various neurodegenerative diseases. But to date, no tests have been able to measure iron in living brain.

Alan Wilman and Gregg Blevins, co-principal investigators from the Faculty of Medicine & Dentistry, used a new MRI method to quantitatively measure iron in the brain to gain a better understanding of what the disease is doing in the brains of those who were recently diagnosed with MS. Twenty-two people with MS took part in the study, along with 22 people who did not have the condition. “In MS, there is a real desire and need to get a good idea of the state and progression of the disease,” says Blevins, who is both a practicing neurologist and a researcher from the Division of Neurology.

“When patients with MS currently get an MRI, the typical measures we look at may not give us a good idea of the nature and state of MS. Using this new MRI method would give physicians a new way to measure the effectiveness



University of Alberta medical researchers who took part in the study: Peter Seres (standing), co-author, Gregg Blevins (left), and Alan Wilman (right).

of new treatments for patients with MS by watching the impact on iron levels. This opens up the idea of having a new biomarker, a new way of looking at the disease over time, watching the disease, seeing the progression or lack of progression of the disease, a new way to track it.”

Wilman, a researcher and physicist in the Department of Biomedical Engineering, says the new MRI method may be a better gauge for disease progression than strictly looking at the number and frequency of relapses.

“This is a new quantitative marker that gives us more insight into MS. We can get a better handle on where patients are at. In terms of clinical symptoms, they may be fine for quite awhile, then

they have a relapse, then they’re fine for quite awhile. Well, the time when they are actually fine, they may not actually be alright.

The disease may be progressing, but there is just no marker right now that shows that. We think the biomarker we have discovered could be an answer. People in the medical research community are very excited about this discovery, because it could be a new way of looking at the disease.”

The new MRI method, which uses a machine that is 90,000 times the strength of the earth’s magnetic field, will give physicians more detail and information about the impact of MS on the brain, insight that doctors and researchers didn’t have before.